



BOLL KING

AUXILIARY NUTRIENT COMPOUND FOR COTTON

**READ ENTIRE LABEL BEFORE USE.
SHAKE WELL BEFORE USE.**

GUARANTEED ANALYSIS

Total Nitrogen		12.00%
5.05% Ammoniacal Nitrogen		
6.95% Urea Nitrogen		
Available Phosphoric Acid (P2O5)		17.00%
Soluble Potash (K2O)		7.00%
Sulfur (S)	4.75%	
Boron (B)	.05%	
Copper (Cu)	.05%	
Iron (Fe)	.10%	
Manganese (Mn)	.05%	
Zinc (Zn)	.05%	

Derived From: Ammonium Phosphate, Urea, Potassium Thiosulfate, Boric Acid, EDTA Copper, HEDTA Iron, EDTA Manganese and EDTA Zinc

CAUTION:

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. In case of contact with eyes, flush immediately with copious amounts of water for 15 minutes. If irritation persists, consult a physician. Avoid contamination of skin and clothing. Wash with soap and water.

WARRANTY AND CONDITION OF SALE: Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied. Buyer and user, therefore, acknowledges and assumes all risk and liability resulting from handling, storage, and use of this material.

Manufactured By: ProTea Botan US, INC.
1039 Autumn Oaks Circle
Collierville TN 38017

NET CONTENTS: 2.5 Gal / 9.45 liters
26.19lbs / 11.9 Kg

DESCRIPTION

BOLL KING is a proprietary patent pending formulation of major, secondary, and micronutrients sequestered in low molecular weight organic acids, and mannose sugars, and proprietary plant extracts. It is for foliar use on cotton at the reproductive stages for improved blooming and fruiting.

COMPATIBILITY

Product is tank mix compatible with most nutrient solutions and insecticides. Avoid extremes in tank pH when mixing. Always check compatibility with standard quart jar method prior to tank mixing.

APPLICATION

Apply as a light foliar mist to the foliage of 5-20 GPA spray solution. Do not over apply to runoff. Allow 5 hours before sprinkle irrigation.

RATES OF APPLICATION:

General rate of application is 1.5 – 2.5 pints per acre depending on crop size and condition.

- **EARLY SEASON:** Apply 1.5 pints per acre immediately prior to pin head square formation.
- **CROP SET:** Apply 2 – 2.5 pints per acre as cotton begins to bloom prolifically.
- **STRESS:** Do not apply to cotton under drought stress. Application can be effective mitigator of temperature stress if soil moisture is adequate.