



DUROX

AUXILIARY NUTRITIONAL COMPOUND

READ ENTIRE LABEL BEFORE USE.

SHAKE WELL BEFORE USE

AVOID EYE CONTACT

DO NOT INGEST

GUARANTEED ANALYSIS

Total Nitrogen		4.00%
3.75% Ammoniacal Nitrogen		
.25% Urea Nitrogen		
Calcium (Ca)	5.00%	
Magnesium (Mg)	1.00%	
Water Soluble Magnesium	1.00%	
Boron (B)	0.75%	

Derived from: Ammonia Nitrate, Urea, Calcium gluconate, Magnesium Sulfate, and Boric Acid

CAUTION:

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. In case of contact with eyes, flush immediately with copious amounts of water for 15 minutes. If irritation persists, consult a physician. Avoid contamination of skin and clothing. Wash with soap and water.

WARRANTY AND CONDITION OF SALE: Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied. Buyer and user, therefore, acknowledges and assumes all risk and liability resulting from handling, storage, and use of this material.

DESCRIPTION

DUROX is a proprietary patent pending formulation of nutrient sequestered in natural mannose sugar and plant extracts. For use on fruits, berries, and vegetables to improve taste, quality, and durability in shipping.

COMPATIBILITY

DO NOT TANK MIX WITH HIGH PHOSPHATE FERTILIZERS. Where compatibility is not known, always pre-test with standard quart jar method prior to tank additions.

HOW TO APPLY

Apply with any equipment that delivers a fine even mist to the foliage of the crop.

GENERAL RECOMMENDATIONS:

- ❖ **BRAMBLES:** (Blackberry, Dew Berry, Raspberry), Apply 1.5-2.0 quarts per acre when floricanes are 1.5-2.0 feet long and repeat in 2.0 weeks. Make an additional application 7-10 days prior to harvest to improve shipping durability.
- ❖ **TOMATOES:** Apply 2.0 quarts per acre immediately prior to bloom, and repeat at 2 week intervals.
- ❖ **PEPPERS** (all types): Apply 1.5-2.0 quarts per acre in early bloom phase and repeat in 2 weeks.
- ❖ **CELERY:** Apply 1.5 quarts per acre at mid stalk development to prevent black heart.
- ❖ **BUSH BERRIES:** (blueberries, huckleberries, goji berries, Saskatoon berries, etc.) For fruit integrity and durability, apply 2.0 qt./acre, (4.67 l/ha), 5-7 days prior to harvest.

Manufactured By: ProTea Botan US, INC., 1039 Autumn Oaks Circle, Collierville, TN 38017

NET CONTENTS: 2.5 Gal / 9.45 liters

27.12 lbs / 12.3 Kg