



# FARM PRO

## FOR SMALL SEEDED GRAINS

**READ ENTIRE LABEL BEFORE USE.  
SHAKE WELL BEFORE USE**

### GUARANTEED ANALYSIS

Total Nitrogen -----	15.00%
6.67% Ammoniacal Nitrogen	
8.33% Urea Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) -----	15.00%
Soluble Potash (K <sub>2</sub> O) -----	5.00%
Sulfur (S))	5.00%
Copper (Cu)	.05%
Iron (Fe)	.10%
Manganese (Mn)	.05%
Zinc (Zn)	.05%

Derived from: Ammonium phosphate, Mono Potassium Phosphate, Potassium tri poly phosphate, Ammonium Sulfate, Ammonium Thiosulfate, Ammonium poly sulfate, Copper EDTA, Iron HEDTA, Manganese EDTA, and Zinc EDTA

### **CAUTION:**

#### **KEEP OUT OF REACH OF CHILDREN**

Harmful if swallowed. In case of contact with eyes, flush immediately with copious amounts of water for 15 minutes. If irritation persists, consult a physician. Avoid contamination of skin and clothing. Wash with soap and water.

Manufactured By: ProTea Botan US, INC., 1039 Autumn Oaks Circle, Collierville, TN 38017

NET CONTENTS: 2.5 gal (9.45 ltrs)  
23.50 lbs (10.68 Kg)

### DESCRIPTION

**FARM PRO** is a proprietary, patent pending formulation of foliar nutrients sequestered in low molecular weight organic acids and fortified with proprietary plant extracts. It is intended as a crop management tool to maximize the genetic potential of cereal grains.

### COMPATIBILITY

Generally **FARM PRO** will be tank mix compatible with most other nutrient sprays, insecticides, fungicides, and herbicides, but where local water sources are unknown, a standard quart jar test should be done prior to tank mixing.

### METHOD OF APPLCIATION

Apply with any equipment that delivers a fine even mist to the foliage of the crop. Generally spray volumes of 5-12 GPA will be sufficient. May be applied with either ground equipment or aircraft.

### GENERAL CROP RECOMMENDATIONS

**FARM PRO** can be applied at any point during the growing cycle, but maximum yield potential can only be realized when the application is made either immediately prior to or early in the grain development stage.

- ❖ WHEAT: Apply 2-3 pints per acre at or near first joint formation.
- ❖ RICE: Apply 2.5 – 4.0 pints per acre immediately prior to green ring stage.
- ❖ OATS, BARLEY, ETC.: Apply 2-3 pints per acre when heads are beginning to form in the stem.

**WARRANTY AND CONDITION OF SALE:** Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied. Buyer and user, therefore, acknowledges and assumes all risk and liability resulting from handling, storage, and use of this material.