



FORAGE BOOST

for HAY, PASTURES AND FOAGE CROPS

READ ENTIRE LABEL BEFORE USE.

SHAKE WELL BEFORE USE

AVOID EYE CONTACT

GUARANTEED ANALYSIS

| | | |
|---------------------------|-------|-------|
| Total Nitrogen | | 4.00% |
| 3.75% Ammoniacal Nitrogen | | |
| .25% Urea Nitrogen | | |
| Calcium (Ca) | 5.00% | |
| Magnesium (Mg) | 1.00% | |
| Water Soluble Magnesium | 1.00% | |
| Boron (B) | 0.75% | |

Derived from: Ammonia Nitrate, Urea, Calcium gluconate, Magnesium Sulfate, and Boric Acid

CAUTION:

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. In case of contact with eyes, flush immediately with copious amounts of water for 15 minutes. If irritation persists, consult a physician. Avoid contamination of skin and clothing. Wash with soap and water.

WARRANTY AND CONDITION OF SALE: Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied. Buyer and user, therefore, acknowledges and assumes all risk and liability resulting from handling, storage, and use of this material.

Manufactured By: ProTea Botan US, INC., 1039 Autumn Oaks Circle, Collierville, TN 38017

NET CONTENTS: 2.5 Gal / 9.45 liters

28.97 lbs / 13.17 Kg

DESCRIPTION

FORAGE BOOST is an auxiliary nutrient formulation for use as a foliar supplement on all hay crops, pasture, and forage crops. It is fortified with low molecular weight organic acids, and proprietary plant extracts. It can be applied at any time for increased quality, rate of growth, and overall yield, but it is particularly effective for regrowth after cutting or grazing.

COMPATIBILITY

FORAGE BOOST is tank mix compatible with most nitrogen solutions, but should never be mixed with phosphate fertilizers. Where compatibility is unknown, always conduct standard quart jar test prior to tank addition.

HOW TO APPLY

Apply with any equipment that delivers a fine even spray mist to plant foliage. Allow 8-10 hours after application before the reintroduction of cattle.

GENERAL RECOMMENDATIONS:

General rate of application is 1.5-2.0 quarts per acre.

- Early season, 1.5-2.0 quarts per acre
- Regrowth after cutting, 2.0-2.5 quarts per acre
- ❖ SWEET SORGHUM: Apply 2.0-2.5 quarts per acre early in growth period.
- ❖ WINTER RYE and FESCUE: Apply 1.5-2.0 quarts per acre. Can be repeated as grazing dictates.
- ❖ ALFALFA: Apply 2.5-3.0 quarts per acre approximately 7-10 days after each cutting.