



RADIX SP

READ ENTIRE LABEL BEFORE USE.

DO EYE CONTACT

DO NOT INGEST

**WASH CONTAMINATED SKIN AND CLOTHING
IMMEDIATELY.**

GUARANTEED ANALYSIS

8-39-7

Total Nitrogen		8.00%
6.34% Ammoniacal Nitrogen		
.73% Nitrate Nitrogen		
.93% Urea Nitrogen		
Available Phosphoric Acid (P2O5)		39.00%
Soluble Potash (K2O)		7.00%
Boron (B)	.02%	
Copper (Cu)	.05%	
Iron (Fe)	.10%	
Manganese (Mn)	.05%	
Molybdenum (Mo)	.0005%	
Zinc (Zn)	.05%	

Derived from: Urea, Potassium Nitrate, Ammonium Phosphate, Boric Acid, Copper EDTA, Iron HEDTA, Manganese EDTA, Sodium Molybdate, and Zinc EDTA

CAUTION:

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. In case of contact with eyes, flush immediately with copious amounts of water for 15 minutes. If irritation persists, consult a physician. Avoid contamination of skin and clothing. Wash with soap and water.

WARRANTY AND CONDITION OF SALE: Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied. Buyer and user, therefore, acknowledges and assumes all risk and liability resulting from handling, storage, and use of this material.

Manufactured By: ProTea Botan US, INC.
1039 Autumn Oaks Circle
Collierville TN 38017

DESCRIPTION

RADIX-SP is a soluble powder formulation of highly concentrated major and micronutrients. It is fortified with natural plant extracts and low molecular weight organic acids. It is for foliar application to root crops and bulb crops to increase size and uniformity.

COMPATIBILITY

Since local water conditions are unknown, a standard quart jar test is recommended prior to tank mixing with any other product or compound.

APPLICATION

The soluble powder should first be slurried before addition to the spray tank. Add to spray tank only under maximum agitation. Apply with any equipment that delivers a fine even spray to the foliage of the crop. For best results use 12-20 GPA spray volume.

GENERAL RECOMMENDATIONS:

- ❖ GARLIC: Apply 1.5-2.0 lbs per acre just as bulbs begin to swell and upsize.
- ❖ ONIONS: (for bulb size increase)
 - Spring applications: Apply 2.0 lbs per acre approximately 30 days after field establishment, and repeat in 30 days. This has proven to be the most effective in yield increases.
 - Summer application alternative: Apply 2.0 lbs./acre 6-8 weeks prior to harvest.
 - Late season alternative: Apply 2.0 lbs./acre one month prior to harvest.

DO NOT MAKE MORE THAN TWO APPLICATIONS PER SEASON.

- ❖ SWEET POTATOES AND YAMS: Apply 2.0 lbs per acre just as vines begin to form and run.
- ❖ WHITE POTATOES: Apply 1.5-2.0 lbs per acre at hook stage.

NET CONTENTS: 24.0 lbs. (10.91 Kg)