



RAPIDE de RIZ

AUXILIARY NUTRITION FOR RICE

READ ENTIRE LABEL BEFORE USE.

- ❖ SHAKE WELL BEFORE USE.
- ❖ AVOID EYE CONTACT.

GUARANTEED ANALYSIS

10-5-5

Total Nitrogen -----	10.00%
0.40% Ammoniacal Nitrogen	
8.0% Urea Nitrogen	
Available Phosphoric Acid (P2O5)-----	5.00%
Soluble Potash (K2O)-----	5.00%
Boron (B)	.02%
Copper (Cu)	.05%
Manganese (Mn)	.05%
Molybdenum (Mo)	.0005%
Zinc (Zn)	.10%

Derived From: Ammonium Phosphate, Potassium Tripoly phosphate, Urea, Boric Acid, Copper EDTA, Manganese EDTA, Sodium Molybdate, and Zinc EDTA

CAUTION:

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. In case of contact with eyes, flush immediately with copious amounts of water for 15 minutes. If irritation persists, consult a physician. Avoid contamination of skin and clothing. Wash with soap and water.

WARRANTY AND CONDITION OF SALE: Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied. Buyer and user, therefore, acknowledges and assumes all risk and liability resulting from handling, storage, and use of this material.

Manufactured By: ProTea Botan US, INC., 1039 Autumn Oaks Circle, Collierville, TN 38017

NET CONTENTS: 2.5 USG / 9.45 liters

22.75 lbs / 10.34 Kg

DESCRIPTION

RAPIDE de RIZ is a proprietary patent pending formulation of major and micronutrients. It is sequestered in a proprietary low molecular weight organic acids, plant fermentation extracts, and high mol natural sugars. It is used as a foliar cover spray on rice for increased yield and quality.

COMPATIBILITY:

RAPIDE de RIZ is tank mix compatible with most nutrient solutions, insecticides, and herbicides. If tank mix compatibility is in question due to local water sources always pre-test with standard quart jar method prior to tank addition.

HOW TO APPLY:

Apply with any equipment that delivers a fine even spray mist to the foliage of the rice. Best results are obtained with sufficient coverage.

- 2-5 GPA is recommended with aircraft application.
- 12-18 GPA is recommended with ground equipment.

RATE OF APPLICATION:

Apply 1.5-2.0 quarts per acre immediately prior to green ring formation on youngest seedlings. An additional application at 1.0 quart per acre at mid-tillering stage is beneficial.