



# COAX-ASR

## FOR PROFESSIONAL TURF USE

READ ENTIRE LABEL BEFORE USE.  
SHAKE WELL BEFORE USE.

### GUARANTEED ANALYSIS

6.5-2-3

Total Nitrogen	6.50%
6.5% Ammoniacal Nitrogen	
Available Phosphoric Acid (P2O5)	2.00%
Soluble Potash (K2O)	3.00%

F1915

Derived from: Phosphoric Acid, Potassium Phosphate, Ammonium lactate and leather hydrolysate

### **CAUTION:**

#### **KEEP OUT OF REACH OF CHILDREN**

Harmful if swallowed. In case of contact with eyes, flush immediately with copious amounts of water for 15 minutes. If irritation persists, consult a physician. Avoid contamination of skin and clothing. Wash with soap and water.

**WARRANTY AND CONDITION OF SALE:** Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied. Buyer and user, therefore, acknowledges and assumes all risk and liability resulting from handling, storage, and use of this material.

Manufactured By: ProTea Botan US, INC.

1039 Autumn Oaks Circle  
Collierville TN 38017

**NET CONTENTS:** 2.5 USG / 9.45 liters

26.38 lbs / 11.98 Kg

**FOR FLORIDA:**

We recommend that you follow the Golf Course BMP's at: <http://www.flaes.org/pdf/Golf%20Course%20BMP's.pdf>

### **DESCRIPTION:**

COAX-ASR is a proprietary formulation of nutrients sequestered in low molecular weight carbon synergists. It is fortified with natural biostimulants, enzymes, and fermented plant extracts. It is for foliar application to all fine turf grasses and is particularly beneficial under environmental stress conditions that create opportunities for pathogens.

### **COMPATIBILITY:**

COAX-ASR can be tank mixed with a wide array of pesticides, but since the effects of the local water source cannot be predicted a standard quart jar test is recommended prior to tank addition.

### **HOW TO APPLY:**

Apply with any equipment that delivers a fine even mist to the turf. Do not overspray to run off. Allow 4 hours after application before sprinkle irrigating.

### **GENERAL RECOMMENDATIONS:**

Begin applications at the onset of turf stress regardless of causal factors. Repeat on a biweekly basis.

- ❖ Apply 3-5 fl. oz. per 1,000 square feet every two weeks.
- ❖ Extreme stress and disease pressure make initial application of 6.0 fl. oz. per 1,000 square feet and repeat weekly at 3.0 fl. oz. per 1,000 square feet while conditions persist.