



**READ ENTIRE LABEL BEFORE USE.
SHAKE WELL BEFORE USE.**

GUARANTEED ANALYSIS

Manganese (Mn) -----	5.0%
Water Soluble Manganese -----	5.0%
F1915	
Derived from: Manganese Citrate	

CAUTION:

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. In case of contact with eyes, flush immediately with copious amounts of water for 15 minutes. If irritation persists, consult a physician. Avoid contamination of skin and clothing. Wash with soap and water.

WARRANTY AND CONDITION OF SALE: Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied. Buyer and user, therefore, acknowledges and assumes all risk and liability resulting from handling, storage, and use of this material.

Manufactured By: ProTea Botan US, INC.
1039 Autumn Oaks Circle
Collierville TN 38017

**NET CONTENTS: 2.5 USG / 9.45 L
23.4 lbs / 10.64 Kg**

FOR FLORIDA:

We recommend that you follow the Golf Course BMP's at: <http://www.flaes.org/pdf/Golf%20Course%20BMP's.pdf>

HOMBRE

**AUXILIARY MANGANESE FORMULATION
FOR SOIL AND FOLIAGE APPLICATION
PROFESSIONAL TURF USE**

DESCRIPTION

HOMBRE is a chelated manganese compound designed for use where manganese availability to the turf is compromised or deficient. **HOMBRE** is designed to furnish rapid manganese availability to the turf, through soil applications, or through foliar applications at the lower rate ranges.

COMPATIBILITY:

HOMBRE is compatible with a wide range of nutrients, pesticides, and fungicides. Do not mix with phosphate. Always determine tank mix compatibility using the standard quart jar method prior to addition to the tank.

APPLICATION:

Apply as a light mist to entire target area.

- Water in soil application
- Do not apply at higher rates as a foliar supplement.

GENERAL RECOMMENDATIONS:

- ❖ Bare Ground And Dormant Turf: Apply at 2-5 fl. oz. per 1000 f². (60-150 ml/100m²), only when turf is browned out due to dormancy or on bare ground prior to establishment.
- ❖ Soil Correction On Actively Growing Turf: Apply at rate of 0.75 to 1.0 fl. oz. per 1000 f². (20-30 ml/100 m²). Repeat every 4-6 weeks.
- ❖ Foliar Supplementation: Apply 0.5-0.75 fl. oz. per 1000 f². (15-20 ml/100 m²). Repeat as necessary.

DO NOT APPLY TO TURF UNDER HEAVY STRESS.