



# REBOUND

PROFESSIONAL TURF USE  
READ ENTIRE LABEL BEFORE USE  
SHAKE WELL BEFORE USE.

## GUARANTEED ANALYSIS

Magnesium (Mg) -----	0.50%
Water Soluble Magnesium -----	0.50%
Boron (B) -----	0.03%
Sulfur (S) -----	2.35%
Iron (Fe) -----	2.50%
Manganese (Mn) -----	1.75%
Zinc (Zn) -----	1.50%
Molybdenum (Mo) -----	0.001%

Derived from: Magnesium Sulfate, Boric Acid, Ferrous Sulfate heptohydrate, Manganese Sulfate, Zinc Sulfate Monohydrate, and Sodium Molybdate Dihydrate

### CAUTION:

#### KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. In case of contact with eyes, flush immediately with copious amounts of water for 15 minutes. If irritation persists, consult a physician. Avoid contamination of skin and clothing. Wash with soap and water.

**WARRANTY AND CONDITION OF SALE:** Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied. Buyer and user, therefore, acknowledges and assumes all risk and liability resulting from handling, storage, and use of this material.

Manufactured By: ProTea Botan US, INC.  
1039 Autumn Oaks Circle  
Collierville TN 38017

NET CONTENTS: 2.5 USG / 9.45 L  
28.0 lbs / 12.73 Kg

### DESCRIPTION:

**REBOUND** is a proprietary formulation of auxiliary plant nutrients designed for use on intensively managed turf such as golf courses and athletic fields. It contains an array of essential nutrients, antioxidants, amino acids, enzymes, natural extracts from sea plants, and diterpine like compounds. When used as directed, **REBOUND** is very effective in turf establishment and recovery from stress conditions.

### COMPATIBILITY:

Always pre-determine tank mix compatibility with standard quart jar method. **DO NOT MIX WITH PHOSPHATES.**

### APPLICATION:

Apply as a light foliar mist in 30-50 GPA, (285-475 L/Ha). DO NOT WATER IN.

### GENERAL RECOMMENDATIONS:

Begin application within 3 weeks after dormancy break or after daytime temperatures reach 75°F(23°C). Apply 1-1.5 fl. oz. per 1000 f<sup>2</sup> (30-45 ml/100m<sup>2</sup>). Repeat every 10-14 days throughout season. For specific deficiency problems, apply 2-4 fl. oz. per 1000 f<sup>2</sup> (60-120 ml per 100 m<sup>2</sup>). If problem persists, repeat in 7-10 days.

### FOR FLORIDA:

We recommend that you follow the Golf Course BMP's at:  
<http://www.flaes.org/pdf/Golf%20Course%20BMP's.pdf>